




090177e1864db9c8\Approved\Approved On: 02-Apr-2015 13:30



(b) (4)






090177e1864db9ce\Approved\Approved On: 02-Apr-2015 13:30



(b) (4)

090177e1864db9c7\Approved\Approved On: 02-Apr-2015 13:30



(b) (4)

090177e1864db9c9\Approved\Approved On: 02-Apr-2015 13:30



(b) (4)

090177e1864db9cb\Approved\Approved On: 02-Apr-2015 13:30

Pharmacist Information:
Usual dosage: Once daily. See package insert for dosage information.
Prepare suspension at time of dispensing as follows: Tap bottle until powder flows freely. Add 131 mL of water for reconstitution. Insert bottle adapter into neck of bottle. Replace bottle cap. Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension. Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser.
Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride. Each bottle contains 750 mg of Methylphenidate Hydrochloride.
Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP controlled room temperature.]
LB8300 Rev 02 09/14

 Pharmacist: Dispense the enclosed Medication Guide to each patient.

Quillivant XR®
methylphenidate HCl
for extended-release oral suspension

750 mg/150 mL total volume
(When reconstituted with 131 mL of water)

25 mg/5 mL
(5 mg/mL) when reconstituted

 Rx only

Shake liquid well before using. If in powder form, do not use, return to Pharmacist.
Keep bottle tightly closed.
Discard any unused portion of the reconstituted suspension after 120 days.
KEEP OUT OF THE REACH OF CHILDREN
Distributed by: NextWave Pharmaceuticals, Inc.,
A subsidiary of Pfizer Inc., New York, NY 10017
Manufactured by: Tris Pharma, Inc.,
Morristown, NJ 08852

LOT: _____
EXP: _____

IMPRINT AREA reads this way NO VARNISH

 **FPO UPC (NDC/HR) @ 100%**
24478-20525 6

14243600

(b) (4)

090177e1864db9cd\Approved\Approved On: 02-Apr-2015 13:30

Pharmacist Information:
Usual dosage: Once daily. See package insert for dosage information.
Prepare suspension at time of dispensing as follows: Tap bottle until powder flows freely. Add 158 mL of water for reconstitution. Insert bottle adapter into neck of bottle. Replace bottle cap. Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension. Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser.
Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride. Each bottle contains 900 mg of Methylphenidate Hydrochloride.
Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP controlled room temperature.]
LB8218 Rev 02 09/14

 Pharmacist: Dispense the enclosed Medication Guide to each patient.

Quillivant XR®
methylphenidate HCl
for extended-release oral suspension

900 mg/180 mL total volume
(When reconstituted with 158 mL of water)

25 mg/5 mL
(5 mg/mL) when reconstituted

 Rx only

Shake liquid well before using. If in powder form, do not use, return to Pharmacist.
Keep bottle tightly closed.
Discard any unused portion of the reconstituted suspension after 120 days.
KEEP OUT OF THE REACH OF CHILDREN
Distributed by: NextWave Pharmaceuticals, Inc.,
A subsidiary of Pfizer Inc., New York, NY 10017
Manufactured by: Tris Pharma, Inc.,
Morristown, NJ 08852

LOT: _____
EXP: _____

IMPRINT AREA reads this way NO VARNISH

 **FPO UPC (NDC/HR) @ 100%**
24478-21030 4

14243600

(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

MITCHELL V Mathis
07/13/2015